

INTERNAL ROOT RESORPTION

The internal aspect of the root is lined by odontoblasts and predentin, which are unable to resorb the dentin. Internal resorption is caused by chronic pulpal inflammation, a disappearance of odontoblasts and predentin, and pulpal invasion of macrophage-like resorbing cells. Generally asymptomatic, these lesions are commonly stumbled upon during routine radiographic evaluation. Pain may occur if the lesion extends to perforate the crown or root. Resorption is active only when part of the pulp remains vital, therefore, pulp testing can be positive. The lesion can occur anywhere along the root surface, but when coronal enough, can cause a pink discoloration by capillaries in the inflamed pulpal tissue under the non-resorbing enamel. It is also relatively rare, occurring in approximately 2% of patients recalled after luxation injuries.

Treatment requires non-surgical root canal therapy to arrest the resorption. If the resorption has progressed far enough and there is external communication to the PDL, then more elaborate steps may need to be taken to save the tooth. Such steps include clinical crown lengthening, flap and restoration, or orthodontic root extrusion. If none of these treatments are an option, then the tooth may require extraction.



Gunraj et al 1999

All cases are patients of Dr. Pamela Nicoara unless otherwise specified.

ROOT RESORPTION: CAUSES AND TYPES

Root resorption is the “physiologic or pathologic process that causes loss of dentin, cementum or bone.” It is usually the result of trauma, chronic inflammation of the pulp or periodontal ligament, or pressure to the periodontal ligament. Reactions on the root surface can affect the adjacent bone, and resorption of the root by dentinoclasts is very similar to resorption of bone by osteoclasts.

There are two main types of resorption: Internal and External. These will be discussed in more detail on the following panels, but an overview is provided:

1. External
 - a. Surface
 - b. Inflammatory
 - c. Replacement
2. Internal

Each type of resorption is relatively easily recognized when in an initial state, however, if the lesions progress and become large enough to encompass internal or external structures, then diagnosis is less clear, treatment becomes more complicated, and the long term prognosis for the tooth decreases.

Advanced Hopeless
Cervical Resorption



Or as adjacent, is too advanced and requires implant replacement.

This issue of **ProbeTips** will review the types of resorption that can occur, and the treatments available to prevent tooth loss.


Copyright 2026 Dr. Pamela Nicoara

Pamela A Nicoara DDS MSD PLLC

PERIODONTOLOGY IMPLANTOLOGY ORAL MEDICINE

Pamela Nicoara is a Board Certified Periodontist practicing in Everett since 2007. She is a UW Perio graduate, and a transplant from Dallas, Texas.

She is driven to achieve esthetic and predictable outcomes, particularly for anterior implant cases, and is always looking to improve processes and results. You can email her directly below with questions, comments, or suggestions for future newsletters.



3125 Colby Avenue, Suite H
Everett WA 98201
T: 425-374-5380 F: 425-374-5382

www.NICOARaperio.com
doctor@NICOARaperio.com

PROBE TIPS

A QUARTERLY PERIODONTAL NEWSLETTER

BY PAMELA NICOARA DDS MSD

Root Resorption Follow Up



Symptomatic
Cervical Resorption



6 years after Endo
& Flap Repair

VOLUME 19, No. 2

AUGUST 2026

Root Resorption Follow Up

EXTERNAL RESORPTION: CONSIDERATIONS

Resorption of the external root structure can be driven by three main factors: trauma, bacterial byproducts from an infected pulpal system, and excessive pressure on the PDL.

The most common forms of **trauma** are luxation or avulsion. Treatment varies depending on the degree of resorption.

If **bacteria** are the sole reason for continued resorption, proper endodontic debridement and obturation should arrest this process. Bacterially driven resorption is usually slower than traumatic resorption.

Consistent or excessive **pressure** on the PDL also causes root resorption. This is seen regularly when deciduous teeth are resorbed by the normal eruption of permanent teeth. If the path of eruption is poor, then an angled third molar can resorb the distal root of a second molar. Pathological causes include orthodontic tooth movement, or slow growing tumors. Once the offending pressure is relieved, then resorption should cease.

EXTERNAL RESORPTION: TYPES

There are three main types of external root resorption: Surface, Inflammatory, and Replacement.

Surface

Surface resorption is a natural process when minor injury or irritation from trauma occurs to the dentin, cementum or PDL. Just as phagocytes migrate to an area of injury to ingest and dispose of unwanted particulate such as broken-down cells, clastic cells on the root surface are attracted to serve their normal scavenging function. Normal scavenging only occurs for about 2-3 weeks if there is no additional stimulation. Therefore, surface resorption usually arrests itself, remains superficial, and there is no treatment necessary.

Inflammatory

In more severe cases of trauma (such as tooth avulsion) where more intense damage to the root surface occurs and the resorptive cells receive continued stimulation for phagocytosis, inflammatory resorption ensues. This is the most common type of external resorption, and although less commonly the cause, periodontal infection or orthodontic tooth pressure can also lead to resorption either laterally or at the apex respectively. If the lesion is deep enough, or the pulp is also infected, then bacterial by-products from the pulp serve as further stimulation for resorption. This type of resorption occurs most frequently and rapidly in immature teeth.

Inflammatory resorption can be classified by its location on the root surface. Of special mention is **cervical resorption** which appears to follow injury to the cervical attachment apparatus only, and is the most common location of external resorption.

Replacement

If trauma is significant enough to cause loss of viability of the PDL cells, such as avulsion with an extended period of time outside of the mouth, then healing occurs without a periodontal attachment apparatus. If less than 20% of the PDL is involved, resorption may be transient and arrest itself. If greater than 20% is involved, resorption progresses and will result in eventual tooth loss. Alveolar bone replaces the resorbed root, and the root dentin fuses to the bone, known as ankylosis. Resorption can occur more rapidly in younger individuals (8-16 years) compared to older patients (17-39 years).

Treatment is extraction, however for growing patients, especially for anterior esthetic cases, rather than remove the tooth when the lesion is first noted, since bone will replace the root and natural ridge augmentation occurs, the tooth can be left in place for as long as possible. This can buy time until growth has ceased so implants can be placed.

Case 1


Treatment of cervical resorption generally involves debridement and restoration of the resorption cavity. Ideally, the use of glass ionomer is best suited to bond to the dentin. It is also more favorable to the gingival tissues than composite as can be seen at the 1 month and 14 year post-operative photos of this female patient initially treated at age 50.

Root canal treatment was necessary because the lesion extended to the pulp.



Case 2

In this case of inflammatory external resorption, the patient, a 71 year old female, was asymptomatic. The resorption was first noted at the gingival margins of #21 and 28, and further radiographic investigation showed significantly more lesions.



Teeth #21, 22, 27 and 28 all had resorptive lesions. All but tooth #21 required endodontic treatment. Flap reflection allowed access for debridement and isolation of the defects for restoration with Fuji II to minimize the soft tissue irritation from the subgingival restorations.

REFERENCES

**Quintessence Int.* Ne et al. 1999. **OSOMOPORE*. Gunraj. 1999. **Glossary-Contemp Terminology for Endo*. AAE. 1994.

complete references available on request